

Provision of Safe Abortion in Armenia

Ruzanna Abrahamyan M.D. Coordinator of the Gynuity Project on Introduction of Medical Abortion in Armenia


> 11 September, 2009, Tbilisi, Georgia




Demographic Trends, Armenia, 1990-2008




Trends of Outcomes of Pregnancy



Women having at least one abortion in the anamneses ~37%



Using Modern Methods of Contraception by Regions, %





01.03.2008-31.03.2009 YSTU in cooperation with Gynuity Health Projects conducted


Clinical research on medical abortion in compliance with following method:

200 mg Mifepristone, after 48 hours Misoprostol – 800 mcg

Medical Abortion in Armenia

Examined:

450 women of 18-47 years



□ repeated pregnancy □ not married □ first pregnancy

Out 450 examined women

31 women had complications in form of incomplete abortion and progressing pregnancy



Out of them:

15 women got additional dose

6 women received additional dose and due to ineffectiveness surgical abortion was provided

10 women were provided with curettage

Side Effects

- •Pain/cramps
- Bleeding
- Nausea, vomiting, diarrhea
- Fever/chills
- Giddiness, headache

Medical Abortion

- High percentage of successful completion
- Surgical intervention is required in small percentage of cases
- Invasion is not required
- Anesthesia is not required
- Rare serious complications
- Fertility is restored nearly immediately after abortion
- Patients have opportunity for controlling procedure, they can choose confidentiality, independence and naturalness

Medical Abortion is Exceptionally Effective and Safe

